

Cal/Ecotox
Exposure Factors for Mourning Dove (Zenaida macroura)*

Page 1

Endpoint Type	Endpoint Value	Error	Range	Units	Sex	Life Stage	Location	Note	Reference
Age at Fledging, Metamorphosis, Weaning	12-13			d	B	Fledgling	Butte; CA	a	1
Age at Fledging, Metamorphosis, Weaning			11-13	d	NR	Fledgling	AL	b	2
Age at Fledging, Metamorphosis, Weaning	review				NR	NR	CANADA; MEXICO; USA	c	3
Age at Sexual Maturity	year of hatching				B	Juvenile	AZ	d	4
Age at Sexual Maturity	93			d	F	Juvenile	VA	e	5
Age at Sexual Maturity	93			d	F	Yearling	AL	f	6
Age at Sexual Maturity	80			d	M	Yearling	AL	g	6
Body Weight - Mean	125			g	B	Adult	Butte; CA	h	1
Body Weight - Mean	112.7	158.8 SD		g	F	Adult	NM	i	7
Body Weight - Mean	99.3	155.1 SD		g	F	Adult	NM	j	7
Body Weight - Mean	114.6	75.7 SD		g	M	Adult	NM	k	7
Body Weight - Mean	111.4	70.7 SD		g	M	Adult	NM	l	7
Body Weight - Mean	129.6	6.9 SD		g	F	Both Adult and Juv.	Lab	m	8
Body Weight - Mean	140.4	10.5 SD		g	M	Both Adult and Juv.	Lab	n	8
Body Weight - Mean	79.0			g	NR	Fledgling	WA	o	9
Body Weight - Mean	7.04			g	B	Hatchling	NE	p	10
Body Weight - Mean	7.7			g	NR	Hatchling	WA	q	9
Body Weight - Mean	114.3	3.3 SE		g	F	Juvenile	VA	r	5
Body Weight - Mean	108	14.7	91-144	g	NR	NR	Lab	s	11
Clutch or Litter Size	1.8				F	Adult	MN	t	12
Clutch or Litter Size	2				F	Adult	ND	u	13
Clutch or Litter Size	1.8				F	Adult	WA	v	14
Clutch or Litter Size	2			eggs/clutch	F	Adult	AL	w	2
Clutch or Litter Size	review				NR	NR	CANADA; MEXICO; USA	x	3
Clutches or Litters per year	5.85-5.98			nests/year	B	Adult	IA	y	15
Clutches or Litters per year	4.4		3-6	nests/yr	B	Adult	TX	z	16
Clutches or Litters per year	review				NR	NR	CANADA; MEXICO; USA	aa	3
Dietary Composition	waste rice, milo, watergrass, yellow star thistle, Turkey mullein seeds, waste wheat, canary grass, safflower, johnson grass				NR	Adult	Butte; CA	ab	1
Dietary Composition	green foxtail (39.8%), yellow foxtail (21.4%), wheat (19.6%), black bindweed (5.1%), maize (4.7%), proso millet (3.0%), flax (2.0%), field mustard (1.6%), lamb's quarters (1.2%)			%	NR	Adult	Lab	ac	17
Dietary Composition	barley, milo, canary grass, flax, lambs quarter, water grass, wheat, cup grass, watermelon, sunflower				B	Both Adult and Juv.	Imperial; CA	ad	18
Dietary Composition	turkey mullein, buckthorn weed, wheat, California poppy, barley, red maids, Napa thistle, miner's lettuce, prostrate pigweed				B	Both Adult and Juv.	San Luis Obispo; CA	ae	18
Dietary Composition	turkey mullein, buckthorn weed, milk thistle, safflower, red maids, California poppy, chickweed, miner's lettuce				B	Both Adult and Juv.	Butte; CA	af	18

Endpoint Type	Endpoint Value	Error	Range	Units	Sex	Life Stage	Location	Note	Reference
Dietary Composition	safflower, milo, turkey mullein, buckthorn weed, California poppy, rice, sunflower, wheat, corn				B	Both Adult and Juv.	Kern; CA	ag	18
Dietary Composition	review				NR	NR	CANADA; MEXICO; USA	ah	3
Dietary Composition	seeds (96.1%), green vegetation (1.1%), insects (2.8%)			%	NR	NR	NM	ai	19
Dietary Composition	seeds (99.9%), green vegetation (0.1%)			%	NR	NR	NM	aj	19
Dietary Composition	haygrazer (15-96%), wheat (trace-31%), pricklepoppy (trace-19%), panic grass (trace-33%), maize (3-76%), pigweed (trace-14%), sunflower (trace-10%), spurge (trace-17%), wild bean (trace-44%), others (1-6%)			%	NR	NR	OK	ak	20
Dietary Composition	halogeton (22.0% +/- 33.9), pigweed (3.9% +/- 13.5), common vetch (4.1% +/-17.9), Indian ricegrass (26.1% +/- 37.6), collomia(6.7% +/- 21.6), barnyard grass (0.3% +/- 5.0), gromwell (0.4% +/- 3.7), wheat (13.2% +/- 29.9), oats (0.6% +/- 4.2), cottonwood (1.6% +/- 11.6), other species (1.7% +/- 7.9), grit (13.9% +/- 23.9)			%	NR	NR	ID	al	21
Duration of Incubation or Gestation	14-15			d	B	Embryo	Butte; CA	am	1
Duration of Incubation or Gestation	14 d, 16 hr to 15 d				NR	Embryo	Lab	an	22
Duration of Incubation or Gestation	14			d	NR	Embryo	AL	ao	2
Duration of Incubation or Gestation	review				NR	NR	CANADA; MEXICO; USA	ap	3
Fledging or Weaning Rate			1.7-2.0		B	Adult	ID	aq	23
Fledging or Weaning Rate	75%				B	Adult	ID	ar	23
Fledging or Weaning Rate	6.0-6.2				B	Adult	Butte; CA	as	1
Fledging or Weaning Rate	64.5			%	B	Adult	Butte; CA	at	1
Fledging or Weaning Rate	90.3%				B	Adult	WA	au	14
Fledging or Weaning Rate	2.1			young/pair	B	Adult	GA	av	24
Fledging or Weaning Rate	48%				B	Adult	GA	aw	24
Fledging or Weaning Rate	21%			%	B	Adult	PUERTO RICO	ax	25
Fledging or Weaning Rate	37.3			%	B	Adult	MO	ay	26
Fledging or Weaning Rate	1.62				B	Fledgling	ID	az	27
Fledging or Weaning Rate	1.8				B	Fledgling	MN	ba	12
Fledging or Weaning Rate	49.4%				B	Fledgling	MN	bb	12
Fledging or Weaning Rate	0.8		0-1.3		B	Fledgling	WA	bc	14
Fledging or Weaning Rate			1.09-1.21	fledglings/n est	B	Fledgling	MN	bd	28
Fledging or Weaning Rate	6.7			fledglings/f emale	B	Fledgling	TX	be	16
Fledging or Weaning Rate	1.87			fledglings/n est	NR	Fledgling	Kern; CA	bf	29

Endpoint Type	Endpoint Value	Error	Range	Units	Sex	Life Stage	Location	Note	Reference
Fledging or Weaning Rate	1.8			fledglings/nest	NR	Fledgling	IA	bg	15
Fledging or Weaning Rate	review				NR	NR	CANADA; MEXICO; USA	bh	3
Food Ingestion Rate	71	66.6-75.4 95% CI		kcal/day/bird	NR	Adult	Lab	bi	17
Food Ingestion Rate	16.1%		7.8-24.1	%	NR	NR	Lab	bj	30
Foraging Distance	review				NR	NR	CANADA; MEXICO; USA	bk	3
Foraging Distance			3-12	km	NR	NR	OK	bl	20
Growth Rate	see figures				B	Nestling	NE	bm	10
Growth Rate	see figures				B	Nestling	ND	bn	13
Growth Rate	review				NR	NR	CANADA; MEXICO; USA	bo	3
Hatching Success	55.2%			%	B	Adult	Kern; CA	bp	29
Hatching Success	67.1%			%	NR	Hatchling	Kern; CA	bq	29
Hatching Success	226 (both eggs); 56 (one egg)				NR	Hatchling	AL	br	2
Longevity	1.53	0.19 SE		yr	B	Adult	Contra Costa; CA	bs	31
Longevity	review				NR	NR	CANADA; MEXICO; USA	bt	3
Metabolic Rate	1.14	0.46 SD		ml O2/g bw/hr	F	Both Adult and Juv.	Lab	bu	8
Metabolic Rate	1.11	0.29 SD		ml O2/g bw/hr	M	Both Adult and Juv.	Lab	bv	8
Population Density	0 to 5 (+/-8)			nests/100 ha	B	Adult	IA	bw	32
Population Density	0.02			nests/ha	B	Adult	ID	bx	23
Population Density	208				B	Adult	ID	by	27
Population Density	1.8				B	Adult	WA	bz	14
Population Density	3.3		1-9		B	Adult	GA	ca	24
Population Density			1-5 per 2.02 ha		B	Adult	Fresno; Kings; CA	cb	33
Population Density	250-330			pairs/220 acres	B	Adult	IA	cc	15
Population Density			65-100	pairs	B	Adult	MN	cd	28
Population Density	165			nests/81 acres	B	Adult	TX	ce	16
Population Density	9.08 (76)			pairs/square mile	B	Adult	ND	cf	34
Population Density	figures			nests/ha	B	Adult	PUERTO RICO	cg	25
Population Density	21.7			birds/100 mi	NR	NR	Kern; CA	ch	29
Surface Area	241			cm^2	NR	Adult	Lab	ci	35
Surface Area	198			cm^2	NR	Adult	Lab	cj	35
Survival/ Mortality	48.0 %	4.3 SE			B	Adult	Contra Costa; CA	ck	31
Survival/ Mortality	28.2%				B	Adult	MN	cl	36
Survival/ Mortality	0.716	0.651-0.782 95% CI			B	Adult	MO	cm	37
Survival/ Mortality	37.3% (before hatch), 9.2% (after hatch)				B	Embryo; Juvenile	MN	cn	28
Survival/ Mortality	13.7%			%	NR	Hatchling	IA	co	15
Survival/ Mortality	58.5%				B	Juvenile	MN	cp	36
Survival/ Mortality	86%				B	Juvenile	TX	cq	16

Exposure Factors for Mourning Dove (Zenaida macroura)*

Page 4

Endpoint Type	Endpoint Value	Error	Range	Units	Sex	Life Stage	Location	Note	Reference
Survival/ Mortality	0.50				B	Nestling	ID	cr	23
Survival/ Mortality	review				NR	NR	CANADA; MEXICO; USA	cs	3
Time of Fledging or Metamorphosis	15			d	B	Fledgling	AL	ct	38
Time of Mating/ Laying	review				NR	NR	CANADA; MEXICO; USA	cu	3
Time of Migration or Dispersal	September				B	Both Adult and Juv.	Butte; CA	cv	1
Time of Migration or Dispersal	winter				B	Both Adult and Juv.	Merced; Stanislaus; CA	cw	39
Time of Migration or Dispersal	summer				B	Both Adult and Juv.	Merced; Stanislaus; CA	cx	39
Time of Migration or Dispersal	review				NR	NR	CANADA; MEXICO; USA	cy	3
Time of Migration or Dispersal	review				NR	NR	CANADA; MEXICO; USA	cz	3
Time of Molt	review				NR	NR	CANADA; MEXICO; USA	da	3
Time of Nesting	April 20-September 20				B	Adult	NE	db	10
Time of Nesting	latter two-thirds of July				B	Adult	ID	dc	27
Time of Nesting	mid-March to September 20				B	Adult	Butte; CA	dd	1
Time of Nesting	Mar-Sep				B	Adult	Kern; CA	de	29
Time of Nesting	April-October				B	Adult	MN	df	28
Time of Nesting	March-September				B	Adult	TX	dg	16
Water Ingestion Rate	6.9	0.9		% body weight/day	NR	NR	Lab	dh	11

- Notes**
- a age at which young leave nest; N=NR; Gray Lodge State Refuge
 - b N=136 nests; near Auburn
 - c N=NR
 - d age of earliest sexual maturity, based on gonad sizes and breeding behaviors; N=75 males, 66 females; near Tuscon
 - e age at which birds were judged capable of reproduction; N=58 birds; Age=131 days; summer, fall
 - f age at which birds became capable of breeding; N=91 birds; Jul-Oct, Jun-Sep
 - g age at which birds became capable of breeding; N=79 birds; Jul-Oct, Jun-Sep
 - h average body weight; N=NR; Gray Lodge State Refuge
 - i mean body weight; N=9 birds; winter; southeastern area
 - j mean body weight; N=6 birds; summer; southeastern area
 - k mean body weight; N=14 birds; winter; southeastern area
 - l mean body weight; N=13 birds; summer; southeastern area
 - m mean body weight; N=14; Condition=fasted; June; captured at Dundee, IL
 - n mean body weight; N=6; Condition=fasted; June; captured at Dundee, IL
 - o average body weight; N=15 nests; Age=14 days; near Orondo, Douglas County
 - p mean body weight; N=28 nestlings; Age=day of hatch; Fremont and Waterloo areas
 - q average body weight; N=15 nests; Age=0 days (hatch); near Orondo, Douglas County
 - r mean body weight; N=5 birds; Age=131 days; summer, fall
 - s average body weight; N=11 birds; Nov-Dec; captured in Santa Monica Mountains
 - t mean clutch size; N=524 nests; April-September; Prairie Island Nuclear Generating Plant
 - u clutch size in 97.2% of observed nests; N=1203 nests; April-September; J. Clark Salyer National Wildlife Refuge
 - v mean number of eggs laid per nesting attempt; N=123 nests; April-August; Rufus Woods Lake area
 - w average clutch size; N=592 clutches; near Auburn
 - x N=NR
 - y average number of nest attempts by individual pairs; N=1108-1443 nests/year; Lewis, Cass County
 - z mean number of nest attempts made per pair per year; N=7 marked pairs; Texas A&M College campus
 - aa N=NR

Exposure Factors for Mourning Dove (*Zenaida macroura*)*

Page 5

ab	food items in crop contents, in order of importance; N=22 birds; November-December; Gray Lodge State Refuge
ac	percent occurrence in diet by weight; N=120 birds; Aug-Sep; captured in Grand Forks County, ND
ad	principal dietary items in crop contents; N=214 crops; all; Imperial Valley; see paper for seasonal changes in relative amounts of items in diet
ae	principal dietary items in crop contents; N=183 crops; all; inner coast mountains; see paper for seasonal changes in relative amounts of items in diet
af	principal dietary items in crop contents; N=152 crops; all; Sacramento Valley; see paper for seasonal changes in relative amounts of items in diet
ag	principal dietary items in crop contents; N=192 crops; all; southern San Joaquin Valley; see paper for seasonal changes in relative amounts of items in diet
ah	N=NR
ai	percent of diet by volume based on crop analysis; N=38 birds; Jul-Sep; New Mexico State University Ranch; See citation for plant seed species consumed.
aj	percent of diet by volume based on crop analysis; N=42 birds; Oct-Mar; New Mexico State University Ranch; See citation for plant seed species consumed.
ak	percent consumption by weight of 9 most frequently eaten foods (range over 7 months); N=NR; Sep-Mar; Jackson County (elev. 430 m); See citation for montly consumption of food items.
al	percent occurrence by mass in crop contents (mean +/- SD); N=223 birds; Jun-Jul; Idaho National Engineering Laboratory
am	incubation period; N=NR; Gray Lodge State Refuge
an	incubation period up to time of hatching; N=1 egg
ao	N=5 cases; near Auburn
ap	N=NR
aq	mean number of fledglings per successful nest (over three years); N=5-7 nests/yr; June-July; Idaho National Engineering Laboratory
ar	proportion of active nests that fledged young; N=24 nests; June-July; Idaho National Engineering Laboratory
as	mean number of nestlings raised per nesting pair per season (representing multiple broods per season); N=43 pairs; Gray Lodge State Refuge
at	proportion of attempted nests that fledged at least one young; N=220 nests; Gray Lodge State Refuge
au	proportion of young fledged in successful nests; N=123 nests; April-August; Rufus Woods Lake area
av	mean productivity; N=73 nests; January-September; southwest and northeast areas
aw	proportion of nests that were successful; N=73 nests; January-September; southwest and northeast areas
ax	nest success estimate based on Mayfield method; N=27 nests; January-December; near Cidra, Guanica, and Cabo Rojo; Nests were considered successful if at least one nestling reached 12 d of age (fledging age).
ay	percent of all nests in which nestlings reached 10 d of age; N=83 nests; Davisdale Wildlife Area (39deg01.3'N,92deg37.5'W); Success in edge habitats did not differ from that in continuous habitats.
az	number of doves fledged per breeding adult; N=NR; June-August; near American Falls
ba	mean number of young fledged per successful nest; N=358 nests; April-September; Prairie Island Nuclear Generating Plant
bb	proportion of active nests that successfully fledged young; N=524 nests; April-September; Prairie Island Nuclear Generating Plant
bc	mean number of young fledged per nesting attempt; N=123 nests; April-August; Rufus Woods Lake area
bd	number of young fledged per nesting attempt; N=81 nests/year; Madelia Research Center, Watonwan County; 61.8% to 67.8% of active nests fledged at least one young.
be	mean number of fledglings produced per adult female; N=7 marked pairs; Texas A&M College campus; Author notes that this value is probably unusually high.
bf	average number of fledglings per successful nest; N=185 successful nests; near Bakersfield; See citation for data from Iowa and Nebraska.
bg	average number of fledglings produced per successful nest; N=1108-1443 nests/year; Lewis, Cass County
bh	N=NR
bi	intake based on analysis of crop contents; N=120 birds; Aug-Sep; captured in Grand Forks County, ND; Equivalent to 16.09 +/- 0.49 g/day/bird (based on a seed diet).
bj	daily food ingestion as percent of body weight; N=22 birds; Indirect measurement, via estimation of loss of body weight over 24 hours.
bk	N=NR
bl	distance flown from night roosts to feeding sites; N=NR; Jun-Dec; Jackson County (elev. 430 m)
bm	growth rates of body weight, body length, wing, alar tract from hatching to fledging; N=53 nestlings; Fremont and Waterloo areas
bn	growth rates of nestling weight, body length, wing chord, tail length, 6th primary; N=178 birds; April-September; J. Clark Salyer National Wildlife Refuge
bo	N=NR
bp	percent of nests in which at least one egg hatched; N=335 nest attempts; near Bakersfield; See citation for data from Iowa and Nebraska.
bq	percent of eggs laid that hatched; N=NR; near Bakersfield; See citation for data from Iowa and Nebraska.
br	number of nests in which one or both eggs hatched; N=592 nests; near Auburn
bs	mean life span; N=725 birds; Berkeley
bt	N=NR
bu	mean oxygen consumption during the light period at 10 deg C; N=14; Condition=fasted; June; captured at Dundee, IL
bv	mean oxygen consumption during the light period at 10 deg C; N=6; Condition=fasted; June; captured at Dundee, IL
bw	mean nest densities in untilled corn or soybean fields; N=8-9 fields; spring; Adair County
bx	density of nests; N=15 plots; June-July; Idaho National Engineering Laboratory

by	number of nests found in 13.4 acres of orchards; N=NR; June-August; near American Falls
bz	nesting pairs per 100 trees checked; N=4 orchards; April-August; Rufus Woods Lake area
ca	mean breeding density; N=ten 100 ac plots; January-September; southwest and northeast areas
cb	range of nesting densities on agroforestry plots; N=6 plots; February-June; San Joaquin Valley Agroforestry Demonstration Project
cc	density of breeding pairs; N=220 acres; Lewis, Cass County
cd	range of total annual nesting population, over 3 yrs; N=65-100 pairs; Madelia Research Center, Watonwan County
ce	peak monthly nesting density in study area; N=648 nests; June; Texas A&M College campus
cf	mean density of breeding birds (maximum density in parentheses); N=130 sample units (160 acres/unit); April-July
cg	Figures of yearly and monthly nest density estimates; N=48-158 transects; January-December; near Guanica and Cabo Rojo
ch	average number of birds per 100 miles of survey route; N=1000 mi surveyed/mo; near Bakersfield; See citation for data from Iowa and Nebraska.
ci	measured area of skin beneath feathers of 649.1 g bird, excluding beak and non-feathered portions of legs; N=1
cj	measured area of the external plumage surface of 649.1 g bird, excluding beak and non-feathered portions of legs; N=1
ck	mean annual mortality rate; N=725 birds; Berkeley
cl	weighted mean annual mortality of banded doves; N=991 birds; near Savage
cm	overall survival of radiomarked individuals; N=315; spring, summer (1 yr); Davisdale and Mussel Fork Wildlife Areas; No differences found among areas, years, or sexes.
cn	percent of eggs or young lost before and after hatch; N=634 eggs; Madelia Research Center, Watonwan County
co	percent of young hatched that did not survive to fledging; N=3580 hatchlings; Lewis, Cass County
cp	weighted mean annual mortality of banded doves; N=1,179 birds; near Savage
cq	estimated mortality rate in the first year, based on band returns; N=NR; Texas A&M College campus
cr	survival rate from incubation to fledging (26 days); N=24 nests; June-July; Idaho National Engineering Laboratory
cs	N=NR
ct	age at fledging; N=35 broods; east-central area
cu	N=NR
cv	period of departure for fall migration; N=NR; Gray Lodge State Refuge
cw	time of northward movement; N=NR; winter; Turlock area
cx	time of southward migration; N=NR; summer; Turlock area
cy	time of juvenile dispersal; N=NR
cz	time of fall and spring migration; N=NR
da	N=NR
db	period during which active nests were observed; N=121 nests; Fremont and Waterloo areas
dc	time of peak nesting activity; N=NR; June-August; near American Falls
dd	nesting season duration (peaks in July); N=NR; Gray Lodge State Refuge
de	period from beginning to end of nesting; N=335 nest attempts; near Bakersfield; See citation for data from Iowa and Nebraska.
df	period from onset of egg laying through fledging; N=~81 nests/year; Madelia Research Center, Watonwan County
dg	period of active nesting; N=648 nests; Texas A&M College campus
dh	daily ad libitum water consumption of captive birds as percent of body weight; N=11 birds; Nov-Dec; captured in Santa Monica Mountains; Minimum water requirement for maintenance of body weight was determined to be 2.8 +/- 1.0 % body weight/day.

References

1	Cowan, John B. 1952. Life history and productivity of a population of western mourning doves in California. Calif. Fish Game. 38(4):505-521.
2	Pearson, Allen M. and George C. Moore. 1939. Nesting habits of the mourning dove in Alabama. Trans. North Am. Wildl. Conf. 4:468-473.
3	Mirarchi, Ralph E. and Thomas S. Baskett. 1994. Mourning Dove (<i>Zenaida macroura</i>), In: Poole, A., P. Stettenheim and F. Gill, eds. The Birds of North America, No. 117. Philadelphia, PA: The Academy of Natural Sciences; Philadelphia PA: The American Ornithologists' Union, Washington, D.C. p 1-32.
4	Irby, Harold D. and Lytle H. Blankenship. 1966. Breeding behaviors of immature mourning doves. J. Wildl. Manage. 30(3):598-604.
5	Mirarchi, R.E., P.F. Scanlon, F.C. Gwazdauskas and R.L. Kirkpatrick. 1981. Gonadal and hormonal characteristics of juvenile female mourning doves in Virginia. Proc. Annu. Conf. Southeast. Assoc. Fish Wildl. Agencies. 34:415-425.
6	White, Lisa M., Ralph E. Mirarchi and Michael E. Lisano. 1987. Reproductive capability of wild hatching-year mourning doves in Alabama. J. Wildl. Manage. 51(1):204-211.
7	Griffing, James P., Paul E. Sawyer and Charles A. Davis. 1977. Mourning dove body weights in southeastern New Mexico during and after a dry spring and early summer. Southwest. Nat. 21(4):545-547.
8	Ivacic, Donald L. and Ronald F. Labisky. 1973. Metabolic responses of mourning doves to short-term food and temperature stresses in winter. Wilson Bull. 85(2):182-196.
9	Cheney, Susan and Philip W. Cheney. 1967. Growth rate of nestling doves. Murrelet. 48(1):14-16.

10 Holcomb, Larry C. and Michael Jaeger. 1978. Growth and calculation of age in mourning dove nestlings. J. Wildl. Manage. 42(4):843-852.

11 MacMillen, Richard E. 1962. The minimum water requirements of mourning doves. Condor. 64(2):165-166.

12 Faanes, Craig A. 1977. Reproductive success of mourning doves near a nuclear generating plant. Loon. 49(4):201-204.

13 Blockstein, David E. 1989. Crop milk and clutch size in mourning doves. Wilson Bull. 101(1):11-25.

14 Knight, Richard L., Dwight G. Smith, David M. Gaudet and Albert W. Erickson. 1984. Nesting ecology of mourning doves in fruit orchards in north-central Washington. Northwest Sci. 58(3):230-236.

15 McClure, Elliot H. 1942. Mourning dove production in southwestern Iowa. Auk. 59:64-75.

16 Swank, Wendell G. 1955. Nesting and production of the mourning dove in Texas. Ecology. 36(3):495-505.

17 Schmid, William D. 1965. Energy intake of the mourning dove, *Zenaida macroura marginella*. Science. 150:1171-1172.

18 Browning, Bruce M. 1962. Food habits of the mourning dove in California. Calif. Fish Game. (2)91-115.

19 Davis, Charles A. 1974. Mourning dove foods in semi-desert, south-central New Mexico. J. Wildl. Manage. 38(4):941-944.

20 Lewis, James C., John A. Morrison, Victor J. Heller and John W. Ault. 1982. Fall-winter habitat use and food habits of doves in southwestern Oklahoma. Proc. Annu. Conf. Southeast. Assoc. Fish Wildl. Agencies. 36:678-690.

21 Markham, O. Doyle, and Charles H. Trost. 1986. Summer foods of mourning doves in southeastern Idaho. Murrelet. 67:60-62.

22 Graber, Richard R. 1955. Artificial incubation of some non-galliform eggs. Wilson Bull. 67(2):100-109.

23 Howe, Frank P. and Lester D. Flake. 1989. Nesting ecology of mourning doves in a cold desert ecosystem. Wilson Bull. 101(3):467-472.

24 Hopkins, Milton N. and Eugene P. Odum. 1953. Some aspects of the population ecology of breeding mourning doves in Georgia. J. Wildl. Manage. 17(2):132-143.

25 Rivera-Milan, Frank F. 1996. Nest density and success of columbids in Puerto Rico. Condor. 98:100-113.

26 Drobney, Ronald D., John H. Schulz, Steven L. Sheriff, and Wesley J. Fuemmeler. 1998. Mourning dove nesting habitat and nest success in central Missouri. J. Field Ornithol. 69(222):299-305.

27 Fichter, Edson. 1959. Mourning dove production in four Idaho orchards and some possible implications. J. Wildl. Manage. 23(4):438-447.

28 Harris, Stanley W., Marius A. Morse, and William H. Longley. 1963. Nesting and production of the mourning dove in Minnesota. Am. Midl. Nat. 69(1):150-172.

29 McClure, H. Elliott. 1950. An eleven-year summary of mourning dove observations in the west. Trans. North Am. Wildl. Nat. Resour. Conf. 15:335-346.

30 Taber, William B., Jr. 1928. A method to determine the weight of food digested daily by birds. Auk. 45(3):339-341.

31 Leopold, A. Starker, and Mark F. Dedon. 1983. Resident mourning doves in Berkeley, California. J. Wildl. Manage. 47(3):780-789.

32 Basore, Nancy S., Louis B. Best, and James B. Wooley, Jr. 1986. Bird nesting in Iowa no-tillage and tilled cropland. J. Wildl. Manage. 50(1):19-28.

33 Kelly, Troy D., Andrew R. Dyer and David L. Chesemore. 1990. Impact of agroforestry plantations grown with agricultural drainwater on avian abundance in the San Joaquin Valley. Trans. West. Sect. Wildl. Soc. 26:97-103.

34 Stewart, Robert E., and Harold A. Kantrud. 1972. Population estimates of breeding birds in North Dakota. Auk. 89:766-788.

35 Walsberg, Glenn E. and James R. King. 1978. The relationship of the external surface area of birds to skin surface area and body mass. J. Exp. Biol. 76:185-189.

36 Thomforde, Lawrence L. 1972. Migration and mortality of banded mourning doves. Minn. Acad. Sci. J. 38(2/3):72-76.

37 Schulz, John H., Ronald D. Drobney, Steven L. Sheriff, Wesley J. Fuemmeler. 1996. Adult mourning dove survival during spring/summer in northcentral Missouri. J. Wildl. Manage. 60(1):148-154.

38 Hitchcock, Ronald R., and Ralph E. Mirarchi. 1986. Dispersal of recently fledged mourning doves from nest sites. Wilson Bull. 98(4):581-585.

39 Channing, Edward. 1979. Movements of banded mourning doves near Turlock, California. Calif. Fish Game. 65(1):23-35.

*Cal/EPA, OEHHA and the University of California Regents are not responsible for damages of any kind resulting from the use of or reliance on information in this report. Users are encouraged to consult the original data. Updated: February 1999.